

Classical dynamics and stability of collapsing thick shells of matter

G.L. Alberghi^{1,2}‡, R. Casadio¹§, and D. Fazi¹||

¹ Dipartimento di Fisica, Università di Bologna, and I.N.F.N, Sezione di Bologna,
Via Irnerio 46, 40126 Bologna, Italy.

² Dipartimento di Astronomia, Università di Bologna, Via Ranzani 1, 40127 Bologna,
Italy.

Abstract. We study the collapse towards the gravitational radius of a macroscopic spherical thick shell surrounding an inner massive core. This overall electrically neutral *macroshell* is composed by many nested δ -like massive *microshells* which can bear non-zero electric charge, and a possibly non-zero cosmological constant is also included. The dynamics of the shells is described by means of Israel's (Lanczos) junction conditions for singular hypersurfaces and, adopting a *Hartree* (mean field) approach, an effective Hamiltonian for the motion of each microshell is derived which allows to check the stability of the matter composing the macroshell. We end by briefly commenting on the quantum effects which may arise from the extension of our classical treatment to the semiclassical level.

1. Introduction

The dynamics of collapsing bodies is a subject which attracted much attention amongst physicists since the very first formulation of General Relativity. However, the general problem of gravitational collapse is extremely complicated and simplified models have therefore been constructed which allow one to describe the dynamics at least for the simplest configurations (for a review of analytical solutions, see Ref. [1]). In this perspective, a lot of effort has been dedicated to the study of the collapse of thin spherical shells (see, e.g. Refs. [2, 3, 4]), for they represent a quite general and simple model and can be regarded as the starting point for the analysis of more complicated systems. In recent years, attention has been focused on quantum effects that can arise from strong gravitational fields characterizing the final stages of the collapse and several approaches have been attempted (for a very limited list, see, e.g. Refs. [5, 6, 7]).

In particular, a formalism was introduced in Ref. [8] which allows one to study the semiclassical behaviour of the shell by adopting a “minisuperspace” approach [9] to its

‡ alberghi@bo.infn.it

§ casadio@bo.infn.it

|| fazi@bo.infn.it

dynamics very much in the same spirit as that previously employed for the Oppenheimer-Snyder model in Ref. [10]. Yet, many aspects of the collapse still need to be investigated at the classical level, which may alter the quantum (semiclassical) picture in a substantial way. For example, the issue of whether the radial degree of freedom of the shell can be described by a quantum wave function strictly depends on the fact that the matter composing the shell be confined within a very small thickness (i.e. of the order of the *Compton* wavelength of the elementary particles composing the shell) around the shell mean radius, so that the assumption of a coherent state is reliable. We refer to this issue as the “stability” problem for thick shells. In fact, a realistic description of the collapse should involve finite quantities (that is, a shell with finite thickness) whereas, already for the classical dynamics, the very use of standard junction equations [3] requires the validity of the *thin shell limit* (i.e. the shell’s thickness be much smaller than its gravitational radius).

In this paper we address the classical dynamics of a collapsing spherical shell (which we call *macroshell*) of proper mass M with an inner massive core of mass M_c and study the conditions under which the matter composing the shell remains confined around its mean radius. In order to verify this property, we discretize the continuous distribution of matter and consider the macroshell as if composed by a large number N of homogeneous nested spherical δ -like *microshells* [11]. Exploiting the spherical symmetry, we can adopt a Schwarzschild-like coordinate system and the motion of the microshells can be described by the usual “areal” radial coordinates r_i ($i = 1, \dots, N$), ordered so that $r_1 < r_2 < \dots < r_N$. Therefore, the thickness of the macroshell is given by $\delta = (r_N - r_1)$ and its mean radius is $R = (r_N + r_1)/2$. Each microshell will be assumed to have the same proper mass $m_i = m$ and a (possibly zero) electric charge Q_i . We will limit our study outside the gravitational radius of the system, $R_H = 2GM_s/c^2$, where M_s is the total ADM mass, so that the radius of the innermost microshell r_1 will always be greater than R_H .

To summarise, the model will be constructed assuming the following conditions:

- (i) $M_c \gg M$. The mass of the core is much bigger than the macroshell’s, so that the total mass-energy of the system is approximately $M_s \simeq M_c$;
- (ii) $m \ll Nm = M$. Each microshell is much lighter than the macroshell;
- (iii) $(R >) r_1 \gg \delta$. The inner radius of the macroshell (and therefore the radius of each microshell) is much larger than its thickness. This implies that the macroshell thickness also remains much smaller than R_H and this condition will limit the temporal range of our analysis;
- (iv) $\sum_{i=1}^N Q_i = 0$. The macroshell is overall electrically neutral;
- (v) $R(0) \gg R_H$ and $\left. \frac{dR}{dt} \right|_{t=0} \simeq 0$. The collapse starts in the weak field regime. One can therefore take t as the Schwarzschild time, although it will then be more convenient to use the (micro)shell proper times (for a detailed comparison, see the Appendix B

in Ref. [11]).

As mentioned above, the classical equations of motion for a thin shell in General Relativity can be obtained from the standard Israel's junction equations. In the present paper, we shall adopt a minisuperspace approach in which the embedding metrics (interior and exterior with respect to the shell) are held fixed and chosen to be particular solutions of the (bulk) Einstein equations, and derive the shell's dynamical equations from a variational principle applied to an effective action. This (equivalent) treatment will allow us to take into account the matter internal degrees of freedom and can be extended to the semiclassical level along the lines of Refs. [8, 11]. In fact, the equations thus obtained are equivalent to the junction conditions, with the exception of one more equation which has no counterpart as a junction condition and represents a secondary constraint for the time variation of the gravitational mass of the shell. The relevant dynamical equation is a first order ordinary differential equation for the shell radius which contains ADM parameters [12] (those defining the inner and outer metrics). In order to determine the shell trajectory, one can fix the initial shell radius, but the initial velocity is then uniquely determined *if* the ADM parameters were also fixed. Alternatively (and more sensibly), one can set the initial radius and velocity and adjust the ADM parameters correspondingly.

The task of extending the above analysis to a system of (micro)shells would be extremely involved in either cases. If we choose to set the initial radii and velocities, then the ADM parameters will have to be reset each time two shells cross. Alternatively, if we set the ADM parameters once and for all, it is possible that the system runs into some spurious (cusp or “superficial”) singularity when two shells cross [1, 13]. In light of a future extension to the semiclassical level, we shall therefore find it more convenient to derive an effective Hamiltonian for the motion of a single microshell in the mean gravitational field generated by the core and all the other microshells treated as a whole. This will allow us to obtain a one-particle effective potential which describes the tidal forces the microshell is subject to, and, in particular, we will see that the binding potential varies with time and confines the microshells within an ever decreasing thickness (at least sufficiently away from the gravitational radius of the system).

We will work in units in which the speed of light $c = 1$ but will explicitly display the gravitational (Newton) constant G .

2. Single shell dynamics

We begin by reviewing the case of one δ -like shell discontinuity. The action for a generic matter-gravity system is given by [14]

$$S = S_{EH} + S_M + S_C , \quad (2.1)$$

where S_{EH} is the bulk Einstein-Hilbert action, S_M the matter action and S_C accounts for boundary contributions.




Figure 1. A schematic view of the relevant space-time. The surface at $r = R_B$ serves to bind the collapsing system inside a “box” of finite spatial extension and the limit $R_B \rightarrow \infty$ can be eventually taken.

In order to describe the motion of a shell, let us consider a space-time volume Ω with boundary $\partial\Omega = \cup_k \Sigma_k$, in which every Σ_k represents a smooth three-dimensional submanifold (see Fig. 1). The action takes the more explicit form (see Refs. [5, 6] and, for a thorough discussion, Ref. [15])

$$S = \frac{1}{16\pi G} \int_{\Omega} d^4x \sqrt{-g} \mathcal{R} + \int_{\Omega} d^4x \mathcal{L}_m + \frac{1}{8\pi G} \int_{\cup_k \Sigma_k} d^3x \sqrt{|^{(3)}g|} \mathcal{K} + \frac{1}{8\pi G} \int_{\cap_k \Sigma_k} d^2x \sqrt{|^{(2)}g|} \sinh^{-1}(\eta), \quad (2.2)$$

where g is the determinant of the four-metric $g_{\mu\nu}$, $\mathcal{R}$ the four-dimensional curvature scalar, $\mathcal{K}$ the extrinsic curvature of the 3-dimensional hypersurface Σ_k whose metric has determinant $^{(3)}g$ and η is the scalar product of the unit normals to two (non-smoothly) intersecting boundaries Σ_k and $\Sigma_{k'}$ whose volume element is denoted by $|^{(2)}g|^{1/2}$. Boundary terms have been included so as to obtain the equations of motion by varying the metric inside Ω with vanishing variations on the border $\partial\Omega$ [2, 15].

In our case, we must also consider the three-dimensional surface Σ swept by the time-like sphere of area $4\pi R^2$ which splits the space-time Ω into two regions: Ω_{in} , internal to the shell, and Ω_{out} , external to the shell (where radiation can be included [16]). Of course, there are an initial hypersurface Σ_i and a final hypersurface Σ_f with possible discontinuities at the shell trajectory’s end-points $\Sigma_{i/f} \cap \Sigma$. Adopting a minisuperspace approach, the metrics inside the two regions $\Omega_{\text{in/out}}$ are chosen to be spherically symmetric solutions of the Einstein field equations and can be written in the

form

$$ds^2 = -A dt^2 + A^{-1} dr^2 + r^2 d\Omega^2, \quad (2.3)$$

in which r is the usual areal coordinate and t is the time variable $t_i \leq t \leq t_f$. The most general solution of this kind (without external radiation) is the Reissner-Nordström-(anti-)de Sitter metric with

$$A_{\text{in/out}} = 1 - \frac{2G M_{\text{in/out}}}{r} + \frac{Q_{\text{in/out}}^2}{r^2} \pm \frac{\Lambda_{\text{in/out}}}{3} r^2, \quad (2.4)$$

where $M_{\text{in/out}}$ is the ADM mass (proper mass plus gravitational energy) enclosed in the sphere of radius r , $Q_{\text{in/out}}$ represents the electric charge inside the sphere of radius r , $\Lambda_{\text{in/out}}$ the cosmological constant (the signs before the last term account for both the de Sitter and the anti-de Sitter metrics).

In Ref. [16], it was shown that, upon choosing a suitable ADM foliation [12] of the space-time, the shell's trajectory can be expressed as $r = R(\tau)$, where τ is an arbitrary time variable on the shell's world volume Σ , and the three metric on Σ is given by

$$ds_\Sigma^2 = -N^2 d\tau^2 + R^2 d\Omega^2, \quad (2.5)$$

where $N = N(\tau)$ is the *lapse function* of the shell. On then integrating over space coordinates, the reduced effective action (2.2) for the shell can be written as

$$\begin{aligned} S_{\text{eff}}(N, R, M_{\text{out}}) &= \frac{1}{G} \int_{\tau_i}^{\tau_f} d\tau \left[\beta R - \dot{R} R \tanh^{-1} \left(\frac{\dot{R}}{\beta} \right) \right]_{\text{out}}^{\text{in}} \\ &\quad - \int_{\tau_i}^{\tau_f} d\tau \left[4\pi N R^2 \rho - \frac{\dot{M}_{\text{out}} R}{2 A_{\text{out}}} + \frac{M_{\text{out}} \dot{R}}{2 A_{\text{out}}} \right] \\ &\equiv \int_{\tau_i}^{\tau_f} d\tau L_{\text{eff}}(N, R, M_{\text{out}}), \end{aligned} \quad (2.6)$$

where $\beta \equiv \sqrt{A N^2 + \dot{R}^2}$, the subscripts “in” and “out” now denote quantities calculated respectively at $r = R - \varepsilon$ and $r = R + \varepsilon$ in the limit $\varepsilon \rightarrow 0^+$, $\rho = \rho(R, M_{\text{out}})$ is the mass-energy density of the shell, and we have introduced the notation $[F]_{\text{out}}^{\text{in}} \equiv F_{\text{in}} - F_{\text{out}}$.

We point out that the action (2.6) allows for the time variation of the ADM mass M_{out} which is a canonical variable of the system, and this corresponds to a radiating shell. In this case, the region Ω_{out} would be further divided into two regions, one enclosing all the radiation and one (Ω_∞ in Fig. 1) devoid of both matter and radiation. In the following Sections, the ADM mass calculated on the external surface of the shell is constant and therefore the last two terms in the action will be dynamically irrelevant. However, we will give the more general expressions for the radiating case below.

Upon varying the action with respect to the canonical variables N , R and M_{out} one obtains the Euler-Lagrange equations of motion for the shell in the form

$$\frac{\delta L_{\text{eff}}}{\delta \vec{X}} \equiv \frac{\partial L_{\text{eff}}}{\partial \vec{X}} - \frac{d}{d\tau} \left(\frac{\partial L_{\text{eff}}}{\partial \dot{\vec{X}}} \right) = 0, \quad (2.7)$$

where $\vec{X} = (N, R, M_{\text{out}})$ and $\delta\vec{X}(\tau_i) = \delta\vec{X}(\tau_f) = 0$. Introducing the canonical momenta conjugated to the variables N, R, M_{out} one obtains

$$P_N \equiv \frac{\partial L_{\text{eff}}}{\partial \dot{N}} = 0 \quad (2.8)$$

$$P_R \equiv \frac{\partial L_{\text{eff}}}{\partial \dot{R}} = - \left\{ \frac{R}{G} \left[\tanh^{-1} \left(\frac{\dot{R}}{\beta} \right) \right]_{\text{out}}^{\text{in}} + \frac{M_{\text{out}}}{2 A_{\text{out}}} \right\} \quad (2.9)$$

$$P_M \equiv \frac{\partial L_{\text{eff}}}{\partial \dot{M}_{\text{out}}} = \frac{R}{2 A_{\text{out}}} . \quad (2.10)$$

We see from the first equation that N is a Lagrange multiplier, corresponding to the effective action being invariant under time reparameterization on the shell. Varying the effective action with respect to N will hence yield a primary constraint instead of a real equation of motion. Eqs. (2.7) thus read

$$\frac{\delta L_{\text{eff}}}{\delta N} = \frac{R}{G N} [\beta]_{\text{out}}^{\text{in}} - 4\pi R^2 \rho = 0 \quad (2.11)$$

$$\begin{aligned} \frac{\delta L_{\text{eff}}}{\delta R} &= \frac{1}{G} \left[\beta + \frac{1}{\beta} \left(R\ddot{R} + \frac{N^2 M}{R} - \frac{R\dot{N}\dot{R}}{N} \right) \right]_{\text{out}}^{\text{in}} + \frac{\dot{M}_{\text{out}}}{\beta_{\text{out}}} \frac{\beta_{\text{out}} - \dot{R}}{1 - 2 M_{\text{out}}/R} \\ &\quad - 4\pi N \frac{\delta(R^2 \rho)}{\delta R} = 0 \end{aligned} \quad (2.12)$$

$$\frac{\delta L_{\text{eff}}}{\delta M_{\text{out}}} = 4\pi N R^2 \frac{\delta \rho}{\delta M_{\text{out}}} - \frac{\beta_{\text{out}} - \dot{R}}{(1 - 2 G M_{\text{out}}/R)} = 0 . \quad (2.13)$$

At this point, one can introduce the Hamiltonian of the system,

$$H = P_R \dot{R} + P_M \dot{M}_{\text{out}} - L_{\text{eff}} = 4\pi N R^2 \rho - \frac{R}{G} [\beta]_{\text{out}}^{\text{in}} , \quad (2.14)$$

and it becomes clear that Eq. (2.11) represents the (secondary) Hamiltonian constraint in the form

$$\frac{H}{N} = 0 , \quad (2.15)$$

whereas Eq. (2.12) can be written as the time preservation of the constraint (2.15),

$$\frac{d}{d\tau} \left(\frac{H}{N} \right) = 0 . \quad (2.16)$$

Making use of the invariance under time reparameterization on the shell, it is convenient to choose the *proper time* gauge $N = 1$, so that the dynamical evolution of the shell is completely determined by the equation

$$H = \frac{R}{G} [\beta]_{\text{out}}^{\text{in}} - 4\pi R^2 \rho = 0 \quad (2.17)$$

and τ is the proper time on the shell. If we consider a shell with constant ADM mass M_{out} , it seems obvious to identify the physical shell energy with its proper mass and set $4\pi R^2 \rho = M$. With this substitution, Eq. (2.17) reads

$$M = \frac{R}{G} \left(\sqrt{A_{\text{in}} + \dot{R}^2} - \sqrt{A_{\text{out}} + \dot{R}^2} \right) , \quad (2.18)$$

where we have used the explicit expression for β and $\dot{R} = dR/d\tau$ denotes the total derivative with respect to the shell's proper time τ .




Figure 2. Evolution of the macroshell's velocity with the radius for $M_s = M$, $M_c = 0$ and $\Lambda = 1$ (in units with $G = 1$).

2.1. Dynamical equation and initial conditions

On squaring twice the above Eq. (2.18) and rearranging we have

$$\dot{R}^2 = \frac{R^2}{4 M^2 G^2} (A_{\text{in}} - A_{\text{out}})^2 - \frac{1}{2} (A_{\text{in}} + A_{\text{out}}) + \frac{M^2 G^2}{4 R^2}. \quad (2.19)$$

It is then clear that the shell's trajectory will be uniquely specified by the initial condition $R(\tau = 0) = R_0$ if the metric functions $A_{\text{in/out}}$ have fixed parameters. One is therefore given two options:

- a) fix the metric parameters and the initial radius $R(0) = R_0$;
- b) set the initial radius $R(0) = R_0$ and velocity $\dot{R}(0) = \dot{R}_0$, and adjust the metric parameters accordingly.

For example, let us consider a macroshell collapsing in a Schwarzschild-de Sitter background, with metrics

$$A_{\text{in}} = 1 - \frac{2 G M_c}{r} + \frac{\Lambda}{3} r^2 \quad (2.20)$$

$$A_{\text{out}} = 1 - \frac{2 G M_s}{r} + \frac{\Lambda}{3} r^2 ,$$

where M_c is the ADM mass of the core and M_s the total ADM mass of the system ¶. Substituting Eqs. (2.20) into Eq. (2.19), we obtain

$$\dot{R}^2 = \left(\frac{M_s - M_c}{M} \right)^2 - 1 + \frac{G (M_s + M_c)}{R} + \frac{G^2 M^2}{4 R^2} - \frac{\Lambda}{3} R^2 , \quad (2.21)$$

which gives the time evolution of the shell's velocity in terms of its radius $\dot{R}(R(\tau))$. Clearly, (as Fig. 2 also shows) that equation describes the collapse of a shell starting

¶ Note that $M_s - M_c > 0$ is not equal to the shell's proper mass M in general. In fact, it can be smaller than M for bound orbits since it also contains the shell's negative gravitational “potential energy”.

from its maximum radius $R_0 = R_{\max}(M, M_s, M_c, \Lambda)$ with zero initial velocity $\dot{R}_0 = 0$. In order to be able to set the initial velocity independently of the initial radius, one can relax one of the metric parameters, e.g., the total ADM mass M_s , which will therefore be determined as $M_s = M_s(R_0, \dot{R}_0; \Lambda, M, M_c)$ by Eq. (2.21) at $\tau = 0$.

2.2. Generalised dynamics

As previously noted, one cannot set the initial radius and velocity if the metric parameters have already been fixed. This, for example, rules out the possibility of setting $M_s = M_c + M$, $\Lambda = 0$ and have the shell start collapsing with zero velocity at finite radius.

Another possibility is to consider the presence of extra (non-gravitational) forces which certainly come into play in the description of astrophysical objects, and allow for equilibrium configurations of the collapsing matter at finite radius (for example, the radiation pressure for main-sequence stars and the degeneracy pressure of the electrons and, at later stages, of the neutrons for compact stars). A way to model that is to introduce an *ad hoc* force which acts upon the system *before* the collapse begins and manages to keep the radius R fixed (for $\tau < 0$). Such a force will be switched off at $\tau = 0$ and the shell then collapses under the pull of just the gravitational force. One can study this (generalized) dynamics of the shell by adding a constant V_0 to the shell effective Hamiltonian,

$$\begin{aligned} H_{\text{eff}} &= \frac{M}{2} \dot{R}^2 + V(R) + V_0 \\ &= \frac{M}{2} \left[\dot{R}^2 - \left(\frac{M_s - M_c}{M} \right)^2 + 1 - \frac{G(M_s + M_c)}{R} - \frac{G^2 M^2}{4 R^2} + \frac{\Lambda}{3} R^2 + \tilde{f}_0 \right], \end{aligned} \quad (2.22)$$

where

$$\begin{aligned} \tilde{f}_0 &= \frac{2}{M} [V(R_0) - V_0] \\ &= \left(\frac{M_s - M_c}{M} \right)^2 - 1 - \frac{G(M_s + M_c)}{R_0} + \frac{G^2 M^2}{4 R_0^2} - \frac{\Lambda}{3} R_0^2, \end{aligned} \quad (2.23)$$

is now an arbitrary constant. We can now set the initial conditions $R(0) = R_0$ and $\dot{R}(0) = 0$ for a given R_0 . Let us finally remark that the Hamiltonian constraint is still given by

$$H_{\text{eff}} = 0, \quad (2.24)$$

and the analysis then proceeds straightforwardly. In fact, Eq. (2.22) for a non-radiating shell is still compatible with the constraint (2.16), since $\tilde{f}_0$ and all the metric parameters are constant in such a case and the dynamical system remains well defined.

3. The macroshell model

In our model, the macroshell is collapsing in a black-hole background and the total ADM mass of the system is constant. In order to study the stability of the matter




Figure 3. Internal structure of the macroshell. We have denoted with $M_< \equiv M_{\text{in}}$ and $M_> \equiv M_{\text{out}}$ the ADM mass calculated respectively on the internal and on the external surface of the microshell of radial coordinate x .

composing the shell, its internal structure must be taken into account and therefore the macroshell is viewed as if composed by many nested microshells. We expect the matter distribution inside the macroshell to alter locally the black-hole background and we study the motion of a single microshell in the mean gravitational field generated by the remaining $(N - 1)$ microshells.

In order to do this, let us single out a microshell of radius x and proper mass $m = M/N$ (see Fig 3). When $x \neq r_i$, we can make use of Eq. (2.17) with $4 \pi x^2 \rho = m$, substitute the expression for β , square twice and rearrange so as to obtain

$$\dot{x}^2 = \frac{x^2}{4 m^2 G^2} (A_< - A_>)^2 - \frac{1}{2} (A_< + A_>) + \frac{m^2 G^2}{4 x^2} , \quad (3.1)$$

where $\dot{x} = dx/d\tau$ denotes the total derivative with respect to the shell's proper time τ . Then, on multiplying Eq. (2.19) by $m/2$, we obtain the equation of motion for a microshell in the form of an effective Hamiltonian constraint, induced by the canonical Hamiltonian constraint (2.17), namely

$$H_{\text{eff}} = \frac{m}{2} \dot{x}^2 + V_{\text{eff}}(x) = 0 , \quad (3.2)$$

and comparison with Eq. (2.19) immediately yields

$$V_{\text{eff}}(x) = \frac{m}{4} (A_< + A_>) - \frac{x^2}{8 m G^2} (A_< - A_>)^2 - \frac{m^3 G^2}{8 x^2} . \quad (3.3)$$

In order to proceed, it would now be necessary to substitute the explicit expressions for $A_<$ and $A_>$, which requires the knowledge of the microshells distribution inside the macroshell. As we discussed in Section 2.1, there are two equivalent ways of determining the evolution of $x = x(\tau)$, one which allows one to set both $x(0) = x_0$ and $\dot{x}(0) = \dot{x}_0$ (with metric parameters correspondingly determined), and one in which the metric parameters are fixed and just $x(0) = x_0$ can be set. However, the former option is presently further complicated by the presence of more than one shell. In fact, whenever




Figure 4. Equivalent configuration in the *thin-shell* limit: (1) $x < X$; (2) $x > X$.

two shells cross (i.e. at the time $\tau = \tau_c$ when $x(\tau_c) = r_i$ for some $i = 1, \dots, N-1$), the metric parameters must be re-evaluated using $x(\tau_c)$ and $\dot{x}(\tau_c)$ as new initial conditions⁺. Since τ_c in turn depends on the trajectory before the crossing, this also means that the system keeps memory of its entire evolution. For this reason, and because we want to be able to extend the treatment to the semiclassical level in the future, we find it more convenient to pursue the other option and fix the metric parameters once and for all. This may imply that the generalised dynamics of Section 2.2 will be used in order to allow us to impose arbitrary initial velocity (e.g. in Section 4). We shall however not display the arbitrary constant $\tilde{f}_0$ in order to keep the expressions simpler.

We then choose the metric parameters according to the assumptions (i)-(v) as outlined in the Introduction. First of all, since $m \ll M \ll M_s$, we shall take $M_> - M_< \simeq m$, which amounts to $\dot{x}(0) \simeq 0$ and is consistent with the choice of having the microshells confined within a thickness δ . With this assumption both $M_<$ and $M_>$ become functions of the microshells distribution $\{(r_i, M_i), i = 1, \dots, N\}$, but, since it is unknown *a priori*, the evaluation of the potential remains involved. A reliable approximation is therefore to adopt a mean field approach and study the motion of the microshell when $x < r_1$ or $x > r_1 + \delta$; in the thin shell limit for the macroshell, all the remaining $(N-1)$ microshells can be viewed as forming a shell of proper mass $(M-m)$ and, if we denote its average radial coordinate by X , we can obtain an equation of motion analogous to Eq. (2.19). In agreement with our gauge choice, we will use the proper times of the (two) shells to foliate the space-time and therefore consider the dynamical variables x and X as functions of the same variable τ . We remark that, if the microshell bears an electric charge Q , then the shell of mass $(M-m)$ must have a charge $-Q$ so that the total macroshell of mass M has zero charge.

One has therefore two distinct situations for $x < X$ and $x > X$ (see cases (1) and

⁺ We do not intend to discuss the issue of shell crossings in any detail here. What happens to two shells when they collide would in fact require the implementation of specific matter properties (such as the detailed treatment of non-gravitational interactions).

(2) in Fig. 4) and two distinct effective potentials:

For $x < X$, the region $R_H < r < x$ has a Schwarzschild-de Sitter metric with ADM mass $(M_s - M)$, the region $x < r < X$ has a Reissner-Nordström-de Sitter metric with ADM mass $(M_s - M + m)$ and electric charge Q , while the outer region $r > X$ has again Schwarzschild-de Sitter metric with ADM mass M_s . We therefore get

$$V_{\text{eff}}^x(x) = \frac{m}{2} \left[-\frac{G}{x} (2M_s - 2M + m) + \frac{Q^2}{Gmx} - \frac{G^2 m^2}{4x^2} + \frac{Q^2}{2x^2} \left(1 - \frac{Q^2}{2G^2 m^2} \right) + \frac{\Lambda}{3} x^2 \right] \quad (3.4)$$

$$V_{\text{eff}}^X(X) = \frac{M-m}{2} \left[-\frac{G}{X} (2M_s - M + m) - \frac{Q^2}{G(M-m)X} - \frac{G^2 (M-m)^2}{4X^2} + \frac{Q^2}{2X^2} \left(1 - \frac{Q^2}{2G^2 (M-m)^2} \right) + \frac{\Lambda}{3} X^2 \right], \quad (3.5)$$

where Eq. (3.4) represents the effective potential for the microshell of proper mass m and radius x , while Eq. (3.5) is the effective potential for the macroshell of proper mass $(M - m)$ and radius X .

The same procedure can be followed for $x > X$. In this case the region $r < X$ has a Schwarzschild-de Sitter metric with ADM mass $(M_s - M)$, the region $X < r < x$ has a Reissner-Nordström-de Sitter metric with ADM mass $(M_s - m)$ and electric charge $-Q$, while the outer region $r > x$ has Schwarzschild-de Sitter metric with ADM mass M_s . This yields

$$V_{\text{eff}}^x(x) = \frac{m}{2} \left[-\frac{G}{x} (2M_s - m) - \frac{Q^2}{Gmx} - \frac{G^2 m^2}{4x^2} + \frac{Q^2}{2x^2} \left(1 - \frac{Q^2}{2G^2 m^2} \right) + \frac{\Lambda}{3} x^2 \right] \quad (3.6)$$

$$V_{\text{eff}}^X(X) = \frac{M-m}{2} \left[-\frac{G}{X} (2M_s - M - m) + \frac{Q^2}{G(M-m)X} - \frac{G^2 (M-m)^2}{4X^2} + \frac{Q^2}{2X^2} \left(1 - \frac{Q^2}{2G^2 (M-m)^2} \right) + \frac{\Lambda}{3} X^2 \right]. \quad (3.7)$$

Since Eq. (3.2) must hold for both the microshell and the macroshell, for each one of the two cases presented above we can write a total effective Hamiltonian for the system which satisfies the constraint

$$\begin{aligned} H_{\text{eff}}^{\text{tot}}(x, X) &\equiv H_{\text{eff}}^x(x) + H_{\text{eff}}^X(X) \\ &\equiv \frac{m}{2} \dot{x}^2 + \frac{M-m}{2} \dot{X}^2 + V_{\text{eff}}^x(x) + V_{\text{eff}}^X(X) \\ &\equiv \frac{m}{2} \dot{x}^2 + \frac{M-m}{2} \dot{X}^2 + V_{\text{eff}}^{\text{tot}}(x, X) = 0. \end{aligned} \quad (3.8)$$

The dynamics has now been reduced to that of a two-body system, so that we can separate the motion of the center of mass from the relative motion by introducing the

center of mass and relative coordinates given by

$$\begin{cases} \bar{r} \equiv x - X \\ R \equiv \frac{m}{M} x + \frac{\mu}{m} X, \end{cases} \quad (3.9)$$

where $\mu \equiv m(M - m)/M$ is the *reduced mass* of the system. The Hamiltonian constraint (3.8) then takes the form

$$H^{(\tau)}(\bar{r}, R) \equiv \frac{\mu}{2} \dot{\bar{r}}^2 + \frac{M}{2} \dot{R}^2 + V^{(\tau)}(\bar{r}, R) = 0, \quad (3.10)$$

where the superscript τ is to recall the time dependence of the Hamiltonian through the variables $\bar{r}$ and R .

Since one component of the system (the macroshell) is much more massive than the other one (the microshell) we have, to lowest order in m/M , $\mu \simeq m$ and we may thus consider the motion of the microshell in the center of mass system to be described by the relative coordinate $\bar{r}$ in much the same way as one treats the motion of the electron in the hydrogen atom. Indeed, since the macroshell is freely falling in the black-hole background, its (mean) radius R and proper time τ define a locally inertial frame. We would thus like to subtract the center of mass motion and obtain a one-particle Hamiltonian for the microshell. In so doing, we obtain an effective Hamiltonian for the whole macroshell of proper mass M and radius R by making use of Eqs. (3.2) and (3.3). In this case we just have two regions, both with Schwarzschild-de Sitter metric, $r < R$ with ADM mass $(M_s - M)$ and $r > R$ with ADM mass M_s ,

$$\begin{aligned} H_M^{(\tau)}(R) &= \frac{M}{2} \dot{R}^2 + V_M^{(\tau)}(R) \\ &= \frac{M}{2} \left[\dot{R}^2 - \frac{2G}{R} \left(M_s - \frac{M}{2} \right) - \frac{M^2 G^2}{4R^2} + \frac{\Lambda}{3} R^2 \right]. \end{aligned} \quad (3.11)$$

Finally, we obtain the one-particle Hamiltonian constraint for the microshell as

$$H_m^{(\tau)}(\bar{r}) \equiv H^{(\tau)}(\bar{r}, R) - H_M^{(\tau)}(R) \simeq \frac{m}{2} \dot{\bar{r}}^2 + V_m^{(\tau)}(\bar{r}) = 0, \quad (3.12)$$

where the potential $V_m^{(\tau)}$ may contain a term $\tilde{f}_0$ so as to ensure that $\bar{r}(0) \simeq \delta/2$ and $\dot{\bar{r}}(0) \simeq 0$, according to the discussion of the generalised dynamics in Section 2.2. Further, the variable R in the potential $V_m^{(\tau)}$ is now to be considered as a (time-dependent) parameter. We note here that this approximation is justified if the typical times of evolution of the macroshell radius R are much larger than the period of a typical oscillation of the microshell around R .

The explicit expression for the potential $V_m^{(\tau)}(\bar{r})$ is extremely complicated and we deem pointless to write it explicitly. Instead, in the next Section, we will plot its behaviour in situations of particular interest.

4. Shell stability

In order to study the stability of the shell we suppose that for $\tau = 0$ all the matter is localized within a thickness $\delta \ll R_H \ll R$ and the shell has negligible (initial) velocity




Figure 5. Effective potential $V_m^{(0)}(\bar{r})/m$ for a macroshell thickness $\delta = 6 \cdot 10^{-9} R_H$ and macroshell radius $R = 10 R_H$. The metric parameters are $M_c = 3 M_\odot$, $M \simeq 8.5 \cdot 10^{-9} M_c \simeq 2.5 \cdot 10^{-8} M_\odot$, $m \simeq 4 \cdot 10^{-9} M \simeq 1 \cdot 10^{-16} M_\odot$, $M_s = M_c + M$, and $\Lambda = Q = 0$, ($M_\odot \simeq 2 \cdot 10^{33} \text{ g}$ is the solar mass and $R_H \simeq 9 \cdot 10^4 \text{ cm}$).

$\dot{R}(0) \simeq 0$. This corresponds to a microshell confined in the region $|\bar{r}| < \delta$ with maximum kinetic energy

$$\frac{m}{2} \dot{r}_{\text{max}}^2 \simeq V_m^{(\tau)}(\bar{r} = \pm\delta) - V_m^{(\tau)}(\bar{r} = 0), \quad (4.1)$$

provided the value $V_m^{(\tau)}(\pm\delta)$ is less than the (possible) maximum of $V_m^{(\tau)}$.

In Fig. 5, we plot the effective potential for a typical case (see the caption for the values of the parameters) at the (relatively) large value of $R = 10 R_H$ and macroshell thickness $\delta \simeq 6 \cdot 10^{-9} R_H$ (we have chosen arbitrary constants so that $V_m^{(0)}(\pm\delta) = 0$). From the graph, it appears that the microshell is classically bound around the “centre of mass” of the macroshell, in agreement with previous results to first order in the small $|\bar{r}|/R$ expansion [11]. This means that if the microshell is initially placed close enough to the macroshell, the mutual common gravitational attraction will keep them together. However, for $|\bar{r}| \gg \delta$, the potential bends down and starts decreasing. This effect cannot be seen in a perturbative expansion for small $|\bar{r}|/R$ but is encoded in the exact potential. One can understand such an effect on considering that if the microshell started with a radius sufficiently larger or smaller than the macroshell’s, the mutual attraction would not be sufficient to overcome the difference in collapsing velocities and their “distance” $|\bar{r}|$ would increase along the collapse. We can note in passing that, as a consequence of the form of the effective potential, one expects that the quantum-mechanical probability for the microshell to tunnel out the well (i.e., from $|\bar{r}| < \delta$ to $|\bar{r}| \gg \delta$) is not zero and (approximately) symmetrical for the cases of a microshell falling inwards or outwards. Strictly speaking, bound states of microshells would therefore be just quantum-mechanically *meta*-stable.




Figure 6. Effective potential $V_m^{(\tau)}/m$ for the same parameters as in Fig. 5 and $R = 10 R_H$ (solid line), $R = 8 R_H$ (dashed line), $R = 6 R_H$ (dotted line), and $R = 4 R_H$ (dash-dotted line).

During the collapse, the velocity of the macroshell $\dot{R}$ increases while its radius R shrinks and the effective potential for the microshell thus changes. This behaviour is easily understood as the increasing of tidal forces between different parts of the macroshell. One might therefore wonder how the time evolution of the binding potential affects the macroshell's thickness and if, eventually, the microshells can escape thus spreading out the system. From the plots in Fig. 6, we can see that the potential $V_m^{(\tau)}$ remains symmetric around the centre but appears to become steeper and steeper as the shell approaches the gravitational radius. In fact, to a given maximum kinetic energy (4.1), there correspond smaller and smaller values of δ for decreasing values of R (moreover, it is easy to see that δ/R actually decreases, since δ decreases faster than R). On the other hand, one may also note that the “width” of the forbidden region decreases (whereas the maxima of the potential increase) and it is thus possible that, at the quantum level, the meta-stable states have different lifetimes. A definite conclusion on this point would need a more complete semiclassical analysis beyond the scope of the present work.

For completeness, we plot the evolving potential when opposite electric charges are carried by the microshell and macroshell in Fig. 7, and when a cosmological constant is included in Fig. 8.

5. Conclusions

We have examined the collapse of a macroscopic self-gravitating spherical shell of matter in the presence of a massive core in the framework of General Relativity. The (macro)shell is assumed to be formed by a large number of thin (micro)shells initially bundled within a very short thickness. The system is then shown to remain classically




Figure 7. Effective potential $V_m^{(\tau)}/m$ for the same parameters as in Fig. 5 except that $Q = 10^{29} e$ (e being the electron's charge) and $R = 10 R_H$ (solid line), $R = 8 R_H$ (dashed line), $R = 6 R_H$ (dotted line), and $R = 4 R_H$ (dash-dotted line).




Figure 8. Effective potential $V_m^{(\tau)}/m$ for the same parameters as in Fig. 5 except that $\Lambda = 10^{-56} \text{ cm}^{-2}$ and $R = 10 R_H$ (solid line), $R = 8 R_H$ (dashed line), $R = 6 R_H$ (dotted line), and $R = 4 R_H$ (dash-dotted line).

stable, since the effective potential which drives the microshells becomes more and more binding as the macroshell approaches the gravitational radius of the system (but remains sufficiently away from it so that our approximations $R - R_H \gg \delta$ and $|\dot{R}| < 1$ hold). Further, including a (reasonably valued) cosmological constant does not alter this behaviour, nor does the presence of electric charges carried by the microshells.

A semiclassical analysis may however change the above conclusion, since the bound states of microshells are just meta-stable and there may be significant changes in their lifetimes (or other, non-adiabatic, effects) as the macroshell collapses. In Refs. [11, 18], it was shown that the microshells can indeed be described by means of localized quantum

mechanical wave functions, however in the linear approximation (first order in $|\bar{r}|/R$) for the effective potential. We have now found that, for sufficiently large values of $|\bar{r}|$ (but still much smaller than R), the complete effective potential reaches maxima and then decreases, thus suggesting a non-vanishing probability for the microshells to tunnel out. It was also found in Refs. [11, 18] that, upon coupling the microshell wave functions to a radiation field, the shell spontaneously emit during the collapse as a non-adiabatic effect. A further analysis of such semiclassical effect in light of the present results is therefore in order, which could also affect the usual thermodynamical description [17].

Let us conclude by mentioning that inclusion of (or a possible relation with) the Hawking radiation [19] as the shell approaches the gravitational radius still remains to be investigated.

References

- [1] R.J. Adler, J.D. Bjorken, P. Chen and J.S. Liu, “Simple analytic models of gravitational collapse,” arXiv:gr-qc/0502040.
- [2] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
- [3] W. Israel, Nuovo Cimento Soc. Ital. Fis., **B 44** (1966) 1; **48** (1966) 463.
- [4] R. Pim and K. Lake, Phys. Rev. D **31** (1985) 233.
- [5] E. Farhi, A.H. Guth and J. Guven, Nucl. Phys. **B 339** (1990) 417.
- [6] S. Ansoldi, A. Aurilia, R. Balbinot, and E. Spallucci, Class. Quantum Grav. **14** (1997) 2727.
- [7] P. Hajicek, Phys. Rev. D **57** (1998) 936; Nucl. Phys. **B 603** (2001) 555.
- [8] G.L. Alberghi, R. Casadio, G.P. Vacca, and G. Venturi, Class. Quantum Grav. **16** (1999) 131.
- [9] B.S. DeWitt, Phys. Rev. **160** (1967) 1113; J.A. Wheeler, in *Batelle rencontres: 1967 lectures in mathematics and physics*, C. DeWitt and J. A. Wheeler editors (Benjamin, New York, 1968).
- [10] R. Casadio and G. Venturi, Class. Quantum Grav. **13** (1996) 2715.
- [11] G.L. Alberghi, R. Casadio, G.P. Vacca, and G. Venturi, Phys. Rev. D **64** (2001) 104012.
- [12] R. Arnowitt, S. Deser and C.W. Misner, *Gravitation: An Introduction to Current Research*, edited by L. Witten (Wiley, New York, 1962).
- [13] T. P. Singh, “Gravitational collapse, black holes and naked singularities,” arXiv: gr-qc/9805066.
- [14] P. Hajicek and J. Kijowski, Phys. Rev. D **57** (1998) 914.
- [15] I. Booth and S. Fairhurst, Class. Quant. Grav. **22** (2005) 4515.
- [16] G.L. Alberghi, R. Casadio, and G. Venturi, Phys. Rev. D **60** (1999) 124018.
- [17] G.L. Alberghi, R. Casadio, and G. Venturi, Phys. Lett. **B 602** (2004) 8; Phys. Lett. **B 557** (2003) 7.
- [18] G.L. Alberghi and R. Casadio, Phys. Lett. B **571** (2003) 245.
- [19] S.W. Hawking, Nature (London) **248** (1974) 30; Commun. Math. Phys. **43** (1975) 199.